

A new approach to estimate directional genetic differentiation and asymmetric migration patterns

Lisa Sundqvist

*University of Gothenburg, Department for Biological and Environmental Sciences,
SE-405 30 Gothenburg, Sweden*

Martin Zackrisson

*University of Gothenburg, Department for Chemistry and Molecular Biology, SE-405 30
Gothenburg, Sweden*

David Kleinhans

*University of Gothenburg, Department for Biological and Environmental Sciences,
SE-405 30 Gothenburg, Sweden; ForWind Center for Wind Energy Research, Institute of
Physics, Carl von Ossietzky University Oldenburg, DE-26129 Oldenburg, Germany*

Abstract

In the field of population genetics measures of genetic differentiation are widely used to gather information on the structure and the amount of gene flow between populations. These indirect measures are based on a number of simplifying assumptions, for instance equal population size and symmetric migration. Structured populations with asymmetric migration patterns, frequently occur in nature and information about directional gene flow would here be of great interest. Nevertheless current measures of genetic differentiation cannot be used in such systems without violating the assumptions. To get information on asymmetric migration patterns from genetic data rather complex models using maximum likelihood or Bayesian approaches generally need to be applied. In such models a large number of parameters are estimated simultaneously and this involves complex optimization algorithms. We here introduce a new approach that intends to fill the gap between the complex approaches and the symmetric measures of genetic differentiation.

Email address: `lisa.sundqvist@bioenv.gu.se` (Lisa Sundqvist)

Our approach makes it possible to calculate a directional component of genetic differentiation at low computational effort using any of the classical measures of genetic differentiation. The approach is based on defining a pool of migrants for any pair of populations and calculating measures for genetic differentiation between the populations and the respective pools. The directional measures of genetic differentiation can further be used to calculate asymmetric migration. The procedure is demonstrated with a simulated data set with known migration pattern. A comparison of the estimation results with the migration pattern used for simulation suggests, that our method captures relevant properties of migration patterns even at low migration frequencies and with few marker loci.

Keywords: Genetic differentiation, Gene flow, Asymmetric Migration, Microsatellite data

1. Introduction

Measures of genetic differentiation are widely used standard tools in population genetics. By measuring the degree of differentiation, information on how populations are structured can be extracted from allelic frequencies. The most frequently used measures are Wright's fixation index F_{st} (Wright, 1943), Nei's G_{st} (Nei, 1973), and the recently introduced D , which is independent of gene diversity (Jost, 2008). In combination with Wright's island model, measures of genetic differentiation can be used to calculate migration between populations (Jost, 2008, Wright, 1931, 1949). The island model is based on a number of simplifying assumptions such as equal population size, drift/migration equilibrium, symmetric migration and no selection or mutation (Whitlock and McCauley, 1999, Wright, 1931). Measures of migration calculated from genetic differentiation share these assumptions, all of which are likely to be violated in natural populations (Whitlock and McCauley, 1999).

Structured populations with asymmetric migration between demes are common in nature, especially in systems driven by physical transport processes such as wind or water currents (Pringle et al., 2011). Both physical processes and density dependent competition can lead to source and sink dynamics in a population and in such populations parameters as genetic differentiation can be significantly skewed (Dias, 1996). In marine environments, ocean currents are known to not only affect the dispersal of planktonic

species, many marine benthic species are also influenced by oceanic circulation since early life-stages (eggs, spores and larvae) often are planktonic (Cowen and Sponaugle, 2009, Siegel et al., 2003). Other examples where gene flow can be affected by physical processes are river systems and wind pollinated plants, moss and lichen (Friedman and Barrett, 2009, Hänfling and Weetman, 2006, Muñoz et al., 2004). In asymmetric systems, it can be essential to estimate directional gene flow to understand the population genetic structure (e.g. Pringle et al., 2011). For instance in conservation, protecting a source population of a threatend species may be more efficient than protecting a sink population. Further all measures of genetic differentiation are symmetric with respect to the populations considered and they do not provide directional information.

To get information on migration in asymmetric systems rather complex models using Maximum likelihood or Bayesian approaches generally need to be applied (e.g. Beerli and Felsenstein, 2001, Wilson and Rannala, 2003). In such models a large number of parameters are estimated simultaneously involving complex optimization algorithms. These models are often used as so called black boxes, implying that users due to the complexity of the approach typically only have a limited knowledge of the underlying model and assumptions.

As a contrast to advanced and computationally expensive methods, we present a rather simple alternative that makes it possible to calculate a directional component of genetic differentiation at low computational effort using any of the classical measures of genetic differentiation. Our approach gives information on direction of gene flow and intends to fill the gap between the complex approaches and the symmetric measures of genetic differentiation. The approach is based on defining a pool of migrants for each pair of populations and calculating measures for genetic differentiation between the populations and the pool. The directional measures of genetic differentiation can further be used to calculate asymmetric migration. This new approach opens up for a new field of applications where directional measures of genetic differentiation can be useful both to explore and to detect asymmetric migration patterns.

2. Measures of genetic differentiation

As a general feature, measures of genetic differentiation are zero if the allele frequencies of the populations are equal (total similarity), values larger

than zero indicate differences in allele frequencies and a value of one indicate no shared alleles (total differentiation) (Meirmans and Hedrick, 2011). Generally they are based on two values, the mean heterozygosity in the total population (H_t) and the mean heterozygosity within the individual populations (H_s) (Meirmans and Hedrick, 2011). For ease of discussion we here introduce vector notations of these parameters and of the measures G_{st} and D .

Let the total number of different alleles present in P populations be N . The allele frequencies in the individual populations can then be arranged in the $N \times P$ -matrix A .

$$A = \begin{pmatrix} a_{11} & \cdots & a_{1P} \\ \vdots & \cdots & \vdots \\ a_{N1} & \cdots & a_{NP} \end{pmatrix} = (\mathbf{a}_1, \cdots, \mathbf{a}_P) \quad , \quad (1)$$

where the matrix element a_{ij} represents the frequency of allele i in population j with $\sum_i a_{ij} = 1$ for any population j . The column vectors $\mathbf{a}_j \in [0, 1]^N$ constitute the allele frequencies in the individual populations.¹ For each population, the degree of heterozygosity (H) can be calculated from the vector of allele frequencies $\mathbf{a} \in [0, 1]^N$ as:

$$H(\mathbf{a}) = 1 - \mathbf{a}^T \mathbf{a} = 1 - |\mathbf{a}|^2 \quad . \quad (2)$$

For a pairwise comparison of two populations with allele frequencies $\mathbf{a} \in [0, 1]^N$ and $\mathbf{b} \in [0, 1]^N$, within-population heterozygosity (H_s) and total-population heterozygosity (H_t) can be calculated from equation (2) as:

$$H_t(\mathbf{a}, \mathbf{b}) = 1 - \frac{1}{4} |\mathbf{a} + \mathbf{b}|^2 \quad , \quad (3a)$$

$$H_s(\mathbf{a}, \mathbf{b}) = 1 - \frac{1}{2} (|\mathbf{a}|^2 + |\mathbf{b}|^2) \quad . \quad (3b)$$

From these expressions the relevant measures for genetic differentiation between populations with allele frequencies $\mathbf{a}$ and $\mathbf{b}$ can be defined using vector algebra:

¹Please note, that bold letters indicate vectors.

$$D_{st}(\mathbf{a}, \mathbf{b}) = H_t(\mathbf{a}, \mathbf{b}) - H_s(\mathbf{a}, \mathbf{b}) = \frac{1}{4}|\mathbf{a} - \mathbf{b}|^2, \quad (4a)$$

$$G_{st}(\mathbf{a}, \mathbf{b}) = \frac{D_{st}(\mathbf{a}, \mathbf{b})}{H_t(\mathbf{a}, \mathbf{b})} = \frac{|\mathbf{a} - \mathbf{b}|^2}{4 - |\mathbf{a} + \mathbf{b}|^2}, \quad (4b)$$

$$D(\mathbf{a}, \mathbf{b}) = \frac{2D_{st}(\mathbf{a}, \mathbf{b})}{1 - H_s(\mathbf{a}, \mathbf{b})} = \frac{|\mathbf{a} - \mathbf{b}|^2}{|\mathbf{a}|^2 + |\mathbf{b}|^2}. \quad (4c)$$

Expressions (4) are equivalent to the (biased) estimators used for calculation of genetic differences as compiled e.g. in (Jost, 2008). A multilocus value of D can be obtained by taking the harmonic mean across loci (Crawford, 2010).

3. A new concept for estimating directional measures of genetic differentiation

Let us start with a thought experiment. Imagine two populations A and B with a strong gene flow from A to B but without any gene flow in the opposite direction, e.g. as observed in systems driven by ocean currents. In addition, population B may be in contact with other populations which are genetically different from A and which contain alleles not present in A. How would such a situation be reflected in the allelic frequencies? Generally we can expect that most alleles present in A are represented in population B, whereas alleles present in B may or may not be present in A due to lack of gene flow from B to A. Hence, in case of neutral mutations, the relative allele frequencies of A should be more or less reflected in B. An idealized example of a allelic matrix is listed in Table 1. In this example allele 3 is present in population B only. Alleles 1 and 2 present in population A are represented at reduced frequencies but equal proportions in population B. From the distribution of allele frequencies it becomes evident, that there is no gene flow from B to A, whereas there might be gene flow from A to B.

How can this concept be formalized? For each pair of populations to be investigated (populations A and B in this example), we introduce a hypothetical *pool of migrants* with an allelic composition inferred from the two populations investigated. That is, we introduce a hypothetical population with an allelic distribution $\mathbf{f}(\mathbf{a}, \mathbf{b})$. From our thought experiment we can define a number of requirements for $\mathbf{f}(\mathbf{a}, \mathbf{b})$:

	Population A	Population B
Allele 1	0.4	0.2
Allele 2	0.6	0.3
Allele 3	0.0	0.5

Table 1: Allelic matrix A of the thought experiment consisting of two populations A and B with directional gene flow from A to B.

1. $\mathbf{f}(\mathbf{a}, \mathbf{b})$ should be symmetric in its arguments, since the order of populations is arbitrary.
2. Alleles not represented in one of the populations can be assumed not to be relevant for migration. As a consequence $\mathbf{f}$ should be non-zero only for alleles present at non-zero frequencies in both populations.

A general form for $\mathbf{f}(\mathbf{a}, \mathbf{b})$ consistent with these two conditions would be

$$f_i(\mathbf{a}, \mathbf{b}) = \sum_{k=1}^K \gamma_k \left(a_i^{\alpha_k} b_i^{\beta_k} + a_i^{\beta_k} b_i^{\alpha_k} \right) \quad \forall i \quad (5)$$

with arbitrary exponents $\alpha_k, \beta_k > 0$ for all k and an arbitrary K defining the number of summands involved. In the optimal example data discussed in connection with the thought experiment (Table 1) we have $a_1/a_2 = b_1/b_2$, which mirrors the fact that alleles 1 and 2 in population B originate from a strong gene flow from A. In order to be consistent with the thought experiment, we require the pool of migrants to reproduce this proportion of alleles as well:

3. In case of strong gene flow in only one direction and genetically totally differentiated source populations we require $f_i(\mathbf{a}, \mathbf{b})/f_j(\mathbf{a}, \mathbf{b}) = a_i/a_j = b_i/b_j$ for all combination of alleles i and j present in both populations.
4. As a vector of allele frequencies $\mathbf{f}$ needs to be normalized, i.e. it needs to fulfill $\sum_i f_i = 1$.

From the 3rd requirement on $\mathbf{f}$ we can demand $\alpha_k + \beta_k = 1$ for any k , making an equation of the form

$$f_i(\mathbf{a}, \mathbf{b}) = \gamma \sum_{k=1}^{\tilde{K}} \tilde{\gamma}_k \left(a_i^{\alpha_k} b_i^{1-\alpha_k} + a_i^{1-\alpha_k} b_i^{\alpha_k} \right) \quad \forall i \quad (6)$$

with arbitrary $0 < \alpha_k < 1$ for all $k \leq \tilde{K}$ the most general functional ansatz for the allelic frequencies in the pooled populations. The constants γ and $\tilde{\gamma}_k$ are related to the previously defined constants γ_k but just as well can be considered as new constants here. For the time being and the rest of this paper we will stick to the simplest available function in this class, namely $\alpha_1 = \tilde{\gamma}_1 = 1/2$ and $\tilde{\gamma}_i = 0 \ \forall i \neq 1$, resulting in

$$f_i(\mathbf{a}, \mathbf{b}) = \gamma \sqrt{a_i b_i} \quad \forall i \quad (7)$$

with $\gamma = (\sum_i \sqrt{a_i b_i})^{-1}$. Thus, $\mathbf{f}(\mathbf{a}, \mathbf{b})$ are the normalized geometrical means of the respective components of $\mathbf{a}$ and $\mathbf{b}$.

As a measure for directional differentiation, we can now compare both A and B respectively to their hypothetical pool of migrants $\mathbf{f}(\mathbf{a}, \mathbf{b})$. For this purpose any of the standard, non-directional measures for genetic differentiation introduced in Section 2 can be applied. With the idealized example data listed in Table 1 we find $\mathbf{f}(\mathbf{a}, \mathbf{b}) = \mathbf{a}$, implying that the genetic constitution of A is equal to the hypothetical pool of migrants, whereas the genetic differentiation between the pool and population B is non-zero ($G_{st}(\mathbf{b}, \mathbf{f}(\mathbf{a}, \mathbf{b})) = 0.19$ and $D(\mathbf{b}, \mathbf{f}(\mathbf{a}, \mathbf{b})) = 0.55$). From this result we can argue that there is very low potential for gene flow from B to A whereas the potential of gene flow from A to B is much higher, which is consistent with the thought example.

In the general case of a number of P populations, we can define a $P \times P$ -matrix B , containing of directional measures for genetic differentiation, as

$$B(A, E(\cdot, \cdot)) = \begin{pmatrix} b_{11}(A, E(\cdot, \cdot)) & \cdots & b_{1P}(A, E(\cdot, \cdot)) \\ \vdots & \cdots & \vdots \\ b_{P1}(A, E(\cdot, \cdot)) & \cdots & b_{PP}(A, E(\cdot, \cdot)) \end{pmatrix} \quad , \quad (8)$$

where the individual elements are defined as

$$b_{ij}(A, E(\cdot, \cdot)) = E(\mathbf{a}_i, \mathbf{f}(\mathbf{a}_i, \mathbf{a}_j)) \quad \forall i, j \quad . \quad (9)$$

Here, $E(\cdot, \cdot)$ is a placeholder for estimates of genetic differentiation, such as G_{st} and D . The value of a particular matrix element b_{ij} can now be understood as an indicator for the potential of migration from population i to population j . Low values indicate high potential for migration and high values indicate low potential for migration in the direction of interest.

4. Estimation of migration

In Wright's Island models F_{st} and also G_{st} can directly be related to migration rates (Wright, 1931). In a similar manner Jost (2008) discusses the dependence of D on migration and mutation and its potential for estimating migration from allelic frequencies (Jost, 2008). If we assume that our data of interest is consistent with the assumptions of the finite island model with infinite number of alleles,² the expression derived in (Jost, 2008) can be used for an estimation of migration rates.

If we assume that the mutation rates are small (i.e. $\mu^2 \ll \mu$) Equation (22) of Jost's 2008 work can be used to estimate the ratio of migration and mutation rate,

$$m/\mu \approx (n-1)(1-D)/D \quad . \quad (10)$$

Since we generally are interested in cases where the migration rate may vary between different pairs of populations, we use Equation (10) with $n = 2$ for an estimation of migration rates between pairs of two populations. We assume, that the mutation rate does not differ between the individual populations. Since we at this stage are interested in the relative sizes of migration rates only, μ can be eliminated from the equation by defining $M = m/\mu$. Relative migration between pairs is then estimated as $M \approx (D-1)/D$.

If we extend this concept to the directional measures of genetic differentiation introduced in the previous section, similar expressions can be used for the estimation of directional, asymmetrical migration rates. For this purpose, we define a $P \times P$ -matrix C of relative migration rates as

$$C(A) = \begin{pmatrix} c_{11}(A) & \cdots & c_{1P}(A) \\ \vdots & \cdots & \vdots \\ c_{P1}(A) & \cdots & c_{PP}(A) \end{pmatrix} \quad , \quad (11)$$

where the individual elements are defined as

$$c_{ij}(A) = \frac{1 - D(\mathbf{a}_i, \mathbf{f}(\mathbf{a}_i, \mathbf{a}_j))}{D(\mathbf{a}_i, \mathbf{f}(\mathbf{a}_i, \mathbf{a}_j))} \quad \forall i, j \quad . \quad (12)$$

²The most important assumptions are: We are dealing with a fixed number of n populations of finite and equal size, interacting through migration of globally dispersing migrants at a certain rate, m . In addition to migration neutral mutations occur at rate μ and result in new alleles. Generally the genetic properties of the model are investigated in equilibrium.

The respective matrix elements $c_{ij}(A)$ here exhibit an estimate for the relative migration rate M from population i to population j and can be evaluated using matrix algebra.

5. Simulations of genetic data with known migration

To test our approach we simulated a system with known migration. The simulation was done in Python (Version 2.7.3) using SimuPop (Version 1.1.0) (Peng and Kimmel, 2005). Allele frequencies were stored in Numpy arrays (Version 1.6.1). Our source code is available as a git-repository at (<https://gitorious.org/ddsim>). Four populations were simulated for 1 500 non-overlapping generations. Each of the populations consists of 1 000 sexually hybrid individuals of a diploid species. Four microsatellite loci on one chromosome were considered and mutation rate was fixed to $\mu = 0.0001$ per allele and generation. The simulation was initialized from allele distributions drawn independently from a pool of 256 alleles. The simulation was run as two phases. First, the populations were evolved without migration for 500 generations. As Figure 1 shows, the number of alleles present in each population drop sharply right after populations are initiated. The argument for this initial, non-migratory phase was to not let genetic drift caused by the artificial population structures of the early generations cloud the results. In the second phase, migration was added to the evolutionary simulation, keeping all other parameters equal.




Figure 1: The number of alleles in the populations over generations

The iterated simulation steps were as follows:

Pre-mating: Mutation A Stepwise Mutator (`StepwiseMutator`) was invoked to simulate microsatellite data. Mutation rate was set to 0.0001 and number of alleles to 256. Default settings were used, with exception

of specifying mutation rate and max allele index as indicated above. With default mutation step size 1, an allele, e.g. number 10 may only evolve into allele number 9 or 11.

Mating: Mating was simulated by random sampling of pairs of parents within each population (`RandomMating`). The two alleles characterizing each offspring was selected randomly, each at a probability of $1 - \mu$ from the respective genetic material of the parents (at equal chance). Default settings were used, except specifying the resulting population sizes (1 000 individuals). After mating the parental generation was removed from the simulation.

Post-mating: Migration (Phase two) During the second phase, migration was invoked via `Migrator`. Migration rates were determined by a migration matrix Mig , all other parameters default. For each i and j a randomly drawn number, based on the probability indicated in Mig_{ij} , of individuals migrate from population i to population j . Individuals could only migrate once and due to the migration, population fluctuate slightly in size around the intended value of 1 000.

The migration matrix Mig used for simulation was:

$$Mig = \begin{pmatrix} 0 & 0.0002 & 0.0001 & 0 \\ 0 & 0 & 0.0002 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (13)$$

Mig represents a migration pattern between the four populations as shown in Figure 2.




Figure 2: The migration pattern as in the matrix Mig .

Results from *our* estimation procedure

At each generation directional D-values (D_d) was calculated for all loci, the harmonic mean of these values over generations is plotted in Figure 3. The estimated matrix elements show strong fluctuations between generations. These stochastic fluctuations as plotted in the figure demonstrates that the accuracy of estimates may differ between generations.




Figure 3: Multilocus D_d -values over generations for the different directions and populations. The figure is divided in two for better visualization of the different lines. All lines are however present in both figures although half of them are transparent.

However the pattern in Figure 3 shows that $1 \rightarrow 2$, $1 \rightarrow 3$ and $2 \rightarrow 3$ generally have lower D_d -values (indication higher migration) then the other dispersal paths. It is also evident that population 4 quite fast reach high differentiation values. At the last generation (1500) a migration matrix (M_{D_d}) was

calculated from the D_d -values according to Equation (12). Since only relative migration is estimated, the migration matrix for clarity was normalized by dividing all elements by the value of the largest matrix element, yielding

$$M_{D_d} = \begin{pmatrix} * & 0.49 & 1.00 & 0.00 \\ 0.01 & * & 0.99 & 0.00 \\ 0.01 & 0.01 & * & 0.00 \\ 0.00 & 0.00 & 0.00 & * \end{pmatrix} . \quad (14)$$

We see that the major patterns from the migration matrix Mig (13) used in the simulation is reproduced, indicating that our approach is able to detect migrations patterns from allele frequency data. Population 4 does not share any alleles with the other three populations and thus shows migration values of zero. The values below the diagonal is low as expected from the pattern shown in Figure 2. Having Figure 2 in mind it is also expected that M_{D_d13} show higher migration then Mig_{13} since population 3 can receive alleles originating from population 1 from two directions. However due to the stochastic fluctuations the accuracy of the method differs between generations and between runs. Since experimentalists typically have access to data from only one generation it is desirable to develop techniques for classifying the accuracy and stability of the estimated migration values, which will be the subject of future studies.

6. Conclusions

We present a new simple and straightforward approach to define a directional component to measures of genetic differentiation. The approach can easily be applied to genetic samples if allele frequencies are known. Directional measures of genetic differentiation can be used for the estimation of asymmetric migration patterns.

7. Acknowledgements

We kindly acknowledge discussions with Per Jonsson, Anna Godhe, Kerstin Johannesson, Bernhard Mehlig. This work was supported by a Linnaeus-grant from the Swedish Research Councils, VR and Formas (<http://www.cemeb.science.gu.se>).

8. References

References

- Beerli, P., Felsenstein, J., 2001. Maximum likelihood estimation of a migration matrix and effective population sizes in n subpopulations by using a coalescent approach. *Proceedings of the National Academy of Sciences* 98 (8), 4563–4568.
- Cowen, R. K., Sponaugle, S., 2009. Larval dispersal and marine population connectivity. *Annual Review of Marine Science* 1, 443–466.
- Crawford, N. G., 2010. Smogd: software for the measurement of genetic diversity. *Molecular Ecology Resources* 10 (3), 556–557.
- Dias, P., 1996. Sources and sinks in population biology. *Trends in Ecology & Evolution* 11 (8), 326–330.
- Friedman, J., Barrett, S. C., 2009. Wind of change: new insights on the ecology and evolution of pollination and mating in wind-pollinated plants. *Annals of Botany* 103 (9), 1515–1527.
- Hänfling, B., Weetman, D., 2006. Concordant genetic estimators of migration reveal anthropogenically enhanced source-sink population structure in the river sculpin, *cottus gobio*. *Genetics* 173 (3), 1487–1501.
- Jost, L., 2008. GST and its relatives do not measure differentiation. *Molecular Ecology* 17 (18), 4015–4026.
- Meirmans, P., Hedrick, P., 2011. Assessing population structure: FST and related measures. *Molecular Ecology Resources* 11 (1), 5–18.
- Muñoz, J., Felicísimo, Á. M., Cabezas, F., Burgaz, A. R., Martínez, I., 2004. Wind as a long-distance dispersal vehicle in the southern hemisphere. *Science* 304 (5674), 1144–1147.
- Nei, M., 1973. Analysis of gene diversity in subdivided populations. *Proceedings of the National Academy of Sciences* 70 (12), 3321–3323.
- Peng, B., Kimmel, M., 2005. simupop: a forward-time population genetics simulation environment. *Bioinformatics* 21 (18), 3686–3687.

- Pringle, J. M., Blakeslee, A. M., Byers, J. E., Roman, J., 2011. Asymmetric dispersal allows an upstream region to control population structure throughout a species range. *Proceedings of the National Academy of Sciences* 108 (37), 15288–15293.
- Siegel, D., Kinlan, B., Gaylord, B., Gaines, S., 2003. Lagrangian descriptions of marine larval dispersion. *Marine Ecology Progress Series* 260, 83–96.
- Whitlock, M. C., McCauley, D. E., 1999. Indirect measures of gene flow and migration: $F_{st} \neq 1/(4nm + 1)$. *Heredity* 82 (2), 117–125.
- Wilson, G., Rannala, B., 2003. Bayesian inference of recent migration rates using multilocus genotypes. *Genetics* 163 (3), 1177–1191.
- Wright, S., 1931. Evolution in mendelian populations. *Genetics* 16 (2), 97.
- Wright, S., 1943. Isolation by distance. *Genetics* 28 (2), 114.
- Wright, S., 1949. The genetical structure of populations. *Annals of Human Genetics* 15 (1), 323–354.